



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Now,

$$\frac{dv}{dr} = \pi h \left[h + (2r - h) \frac{dh}{dr} \right], \quad \text{and} \quad \frac{dh}{dr} = 1 - \csc \alpha.$$

Therefore,

$$\frac{dv}{dr} = \pi h [a \csc \alpha - r(\csc \alpha - 1)(\csc \alpha + 2)].$$

Obviously, $h = 0$ cannot give a maximum overflow. Consequently,

$$r = \frac{a \csc \alpha}{(\csc \alpha - 1)(\csc \alpha + 2)} = \frac{a \sin \alpha}{(1 - \sin \alpha)(1 + 2 \sin \alpha)} = a \sin \alpha / (\sin \alpha + \cos 2\alpha).$$

Furthermore,

$$\frac{d^2v}{dr^2} = -\pi h (\csc \alpha - 1)(\csc \alpha + 2).$$

Since $h \neq 0$ and $\csc \alpha > 1$, d^2v/dr^2 is negative so that the condition for a maximum is satisfied.

Also solved by A. H. WILSON, S. E. RASOR, ELMER SCHUYLER, GEORGE RAYNOR, C. N. SCHMALL, H. L. AGARD, C. N. MILLS, LEWIS CLARK, L. M. PICKETT, C. A. BERGSTRESSER, C. HORNUNG, L. G. WELD, J. A. BULLARD, G. W. HARTWELL, J. V. BALCH, J. A. WHITTED, ELIJAH SWIFT, J. C. RAYWORTH, and J. A. CAPARO.

396. Proposed by ELBERT H. CLARKE, Purdue University.

The length of the curve $y = x^n$ from the origin to the point $(1, 1)$ is given by the formula

$$= \int_0^1 \sqrt{1 + n^2 x^{2n-2}} \cdot dx.$$

Our geometric intuition would tell us that the limit of this length as n becomes infinite is 2. Give a strict analytic proof that

$$\lim_{n \rightarrow \infty} \int_0^1 \sqrt{1 + n^2 x^{2n-2}} \cdot dx = 2.$$

SOLUTION BY ELIJAH SWIFT, University of Vermont.

We can easily show that

$$(1) \quad \lim_{n \rightarrow \infty} n^2 x^{2n-2} = 0, \quad 0 \leq x < 1.$$

Next, we choose a positive number, ϵ , as small as we like. We can then find a value N of n such that for it and all larger values

$$(2) \quad n^2(1 - \epsilon)^{2n-2} < \eta,$$

where η is a positive number as small as we like. This is possible on account of (1).

Let x_n be a value of x such that $n^2 x_n^{2n-2} = 2$.

Now write the given integral, I , as a sum of three integrals.

$$I = \int_0^{1-\epsilon} \sqrt{1 + n^2 x^{2n-2}} \cdot dx + \int_{1-\epsilon}^{x_n} \sqrt{1 + n^2 x^{2n-2}} \cdot dx + \int_{x_n}^1 \sqrt{1 + n^2 x^{2n-2}} \cdot dx.$$

Considering values of $n > N$, we have

$$(3) \quad \left| \int_0^{1-\epsilon} \sqrt{1 + n^2 x^{2n-2}} \cdot dx - 1 \right| = \left| \int_0^{1-\epsilon} \{ \sqrt{1 + n^2 x^{2n-2}} - 1 \} dx - \epsilon \right| \\ \leq \left| \int_0^{1-\epsilon} \{ \sqrt{1 + \eta} - 1 \} dx \right| + \epsilon < \frac{\eta}{2} + \epsilon. \quad (\text{See (2).})$$

This holds if $n \geq N$.

Now

$$(4) \quad \int_{1-\epsilon}^{x_n} \sqrt{1 + n^2 x^{2n-2}} \cdot dx < \int_{1-\epsilon}^1 \sqrt{1 + 2} \cdot dx = \sqrt{3} \cdot \epsilon.$$

Finally, since for values of x greater than x_n , $n^2 x^{2n-2} > 2$, we may develop $\sqrt{1 + n^2 x^{2n-2}}$ into a series in descending powers of $n^2 x^{2n-2}$, and the series will be uniformly and absolutely convergent for values of x between, and including, x_n and 1, and all values of $n \geq N$, we have

$$\sqrt{1 + n^2 x^{2n-2}} = nx^{n-1} + \frac{1}{2} \cdot \frac{1}{nx^{n-1}} - \frac{1}{2 \cdot 4} \cdot \frac{1}{n^3 x^{3n-3}} + \dots$$

Hence,

$$(5) \quad \int_{x_n}^1 \sqrt{1 + n^2 x^{2n-2}} \cdot dx = 1 - x_n^n + \left[\frac{1}{2} \cdot \frac{1}{n(-n+2)} x^{-n+2} - + \dots \right]_{x_n}^1.$$

Since,

$$(6) \quad nx_n^{n-1} = \sqrt{2},$$

$$(7) \quad \lim_{n \rightarrow \infty} x_n^n = 0.$$

We may now obtain the limit of the right hand side of (5) when $n \rightarrow \infty$ by taking the limit of each term. This gives, in consequence of (6) and (7), 1 as the limit. In other words, we may choose n so large that

$$(8) \quad \int_{x_n}^1 \sqrt{1 + n^2 x^{2n-2}} \cdot dx = 1 + \zeta,$$

where ζ is a constant as small numerically as we please for all values of $n > N$.

Then

$$\begin{aligned} \left| \int_0^1 \sqrt{1 + n^2 x^{2n-2}} \cdot dx - 2 \right| &\leq \left| \int_0^{1-\epsilon} \sqrt{1 + n^2 x^{2n-2}} \cdot dx - 1 \right| + \left| \int_{1-\epsilon}^{x_n} \sqrt{1 + n^2 x^{2n-2}} \cdot dx \right| \\ &\quad + \left| \int_{x_n}^1 \sqrt{1 + n^2 x^{2n-2}} \cdot dx - 1 \right| < \frac{\eta}{2} + \epsilon + \sqrt{3} \cdot \epsilon + |\zeta|. \end{aligned}$$

Since this may be made as small as we please by a suitable choice in order of ϵ , η , and ζ , we have proved the statement.

Also solved by TOBIAS DANTZIG.

MECHANICS.

312. Proposed by C. N. SCHMALL, New York City.

A ball of elasticity e is projected upward from a point on an inclined plane, so that after its first contact with the plane it rebounds to its starting point. If ϕ be the inclination of the plane to the horizontal, and ψ the angle made by the line of projection with the inclined plane, show that

$$\cot \phi \cot \psi = e + 1.$$


I. SOLUTION BY HORACE OLSON, Chicago, Illinois.

In order that the answer given may be true the ball must be thrown in a vertical plane perpendicular to the inclined plane. Taking the axis of coordinates as indicated in the figure, the equations of the trajectory are:

$$x = vt \cos \psi - \frac{gt^2 \sin \phi}{2},$$

$$y = vt \sin \psi - \frac{gt^2 \cos \phi}{2},$$

t being the time from the instant of projection. Then,



$$\frac{dx}{dt} = v \cos \psi - gt \sin \phi \quad \text{and} \quad \frac{dy}{dt} = v \sin \psi - gt \cos \phi.$$